

Probability

| Combining Events

Prof. Dr. Mostafa Elhosseini

<https://youtube.com/drmelhosseini>

| Example

- An electrical engineer has on hand **two boxes of resistors**, with **four** resistors in each box. The resistors in the first box are labeled **10 Ω** (ohms), but in fact their resistances are **9, 10, 11, and 12 Ω** . The resistors in the second box are labeled **20 Ω** , but in fact their resistances are **18, 19, 20, and 21 Ω** . The engineer chooses one resistor from each box and determines the resistance of each.
- Let **A** be the event that the **first resistor has a resistance greater than 10**, let **B** be the event that the **second resistor has a resistance less than 19**, and let **C** be the event that the **sum of the resistances is equal to 28**.
- Find a **sample space** for this experiment, and specify the subsets corresponding to the events **A, B, and C**.

| Example

- **S** = {(9, 18), (9, 19), (9, 20), (9, 21), (10, 18), (10, 19), (10, 20), (10, 21), (11, 18), (11, 19), (11, 20), (11, 21), (12, 18), (12, 19), (12, 20), (12, 21)}
- **A** = {(11, 18), (11, 19), (11, 20), (11, 21), (12, 18), (12, 19), (12, 20), (12, 21)}
- **B** = {(9, 18), (10, 18), (11, 18), (12, 18)}
- **C** = {(9, 19), (10, 18)}

| Combining Events

- We often construct events by combining simpler events.
- Because events are subsets of sample spaces, it is traditional to use the notation of sets to describe events constructed in this way

| Union

- The **union of two events A and B**, denoted **$A \cup B$** , is the set of outcomes that belong either to A, to B, or to both.
- In words, $A \cup B$ means “A or B.” Thus the event $A \cup B$ occurs whenever either A or B (or both) occurs.

| Intersection

- The **intersection of two events A and B**, denoted $A \cap B$, is the set of outcomes that belong both to A and to B.
- In words, $A \cap B$ means “A and B.” Thus the event $A \cap B$ occurs whenever both A and B occur.

| Complement

- The **complement of an event A** , denoted A^c , is the set of outcomes that do not belong to A . In words, A^c means “not A .” Thus the event A^c occurs whenever A does not occur.

| Example

- Refer to previous Example. Find $B \cup C$ and $A \cap B^c$.

| Example

- The event $B \cup C$ contains all the outcomes that belong either to B or to C , or to both.
- Therefore $B \cup C = \{(9, 18), (10, 18), (11, 18), (12, 18), (9, 19)\}$
- The event B^c contains those outcomes in the sample space that do not belong to B .
- It follows that the event $A \cap B^c$ contains the outcomes that belong to A and do not belong to B . Therefore
- $A \cap B^c = \{(11, 19), (11, 20), (11, 21), (12, 19), (12, 20), (12, 21)\}$

| Mutually Exclusive Events

- There are some **events that can never occur together**.
- For example, it is impossible that a **coin can come up both heads and tails**, and it is impossible that a steel pin can be both **too long and too short**. Events like this are said to be mutually exclusive
- The events A and B are said to be mutually exclusive if they have **no outcomes in common**.
- More generally, a collection of events **$A_1, A_2, \dots, A_n$** is said to be mutually exclusive if **no two of them have any outcomes in common**.

| Example

- Refer to Previous Example. If the experiment is performed, is it possible for events A and B both to occur? How about B and C? A and C? Which pair of events is mutually exclusive?

| References

- Understanding Boxplots <https://towardsdatascience.com/understanding-boxplots-5e2df7bcbd51>
- What does a box plot tell you?
<https://www.simplypsychology.org/boxplots.html>
- Statistics for Engineers and Scientists, Fourth Edition, William Navidi
Colorado School of Mines
- Khan Academy
<https://www.khanacademy.org/math/cc-sixth-grade-math/cc-6th-data-statistics/cc-6th-box-whisker-plots/v/constructing-a-box-and-whisker-plot>